



Vol. 3 No. 2 (February) (2025)

Nano-Enabled Strategies for Soil Fertility and Plant Productivity: A Sustainable Approach

Asiya Akbar

Department of Agricultural and Environmental Sciences, The Islamia University of Bahawalpur: Email: asiyaakbar413@gmail.com

Aqsa Yaqoob

Postgraduate student, Department of Botany, University of Agriculture Faisalabad, Pakistan. Email: aqsayaqoob555@gmail.com

Fatima Asif

Sustainable development study centre SDSC, Government College University, Lahore Email: fatimaasif1111@gmail.com

Muhammad Waqar

Department of Botany, The Islamia University Bahawalpur.
Email: mwaqarmission@gmail.com

Misbah Rani

Department of Biotechnology, Lahore College for Women University, Lahore.
Email: misbahrani780@gmail.com

Qurat Ul Ain Hyder

Department of Botany, Faculty of Life Sciences, The Women University Multan,
Email: quratulainhyder12@gmail.com

Hafiz Muhammad Usman

Department of Agricultural and Environmental Sciences, The Islamia University of Bahawalpur: Email: hu542652@gmail.com

Abstract

Nanotechnology has recently emerged as another innovative technology that concerns agriculture. It brings solutions to plant growth, soil fertility, and crop yield. Current studies are aimed at assessing the effects of zinc oxide (ZnO) and titanium dioxide (TiO₂) nanoparticles on seed germination, plant growth, biochemical response, and soil microbial activities. Plants were subjected to various concentrations of nanoparticles, and the impacts on the rest of the physiological-biochemical parameters were determined. Hence, it was revealed that seed germination percentage increased chlorophyll content and biomass production, though the differences in ZnO and TiO nanoparticles were significant. The roots were more extensive, and nutrient uptake also was increased because the nanoparticles control the action of auxins and enhance nutrients in the soil. The high antioxidant enzyme activity in biochemical analyses also resulted in reduced oxidative damage and increased resistance of the plant-soil analyses indicating that nitrogen and phosphorus availability was enhanced with a concomitant positive effect on microbial communities at optimal levels of nanoparticles. However, phytotoxic effects were noted at higher



Vol. 3 No. 2 (February) (2025)

concentrations, emphasizing the need for dose optimization. Indeed, statistical analysis validated the finding and indicated that these nanoparticles hold great promise for application as green inputs in agriculture. More studies need to be carried out in the long term on environmental implications and develop safe means of application for large-scale agricultural use since, in this study, ZnO and TiO₂ nanoparticles seem to have a positive influence on plant growth and soil enhancement.

Introduction

Nanotechnology is emerging as a revolutionary force in the present practices in agriculture, soil science, and botany that are providing new avenues towards plant growth stimulation, soil fertility improvement, and sustainable environment-friendly agriculture[1]. Since population increase everywhere is driving up the need for food to be produced at a faster pace, there is indeed the need to carry out resourceful and minimally harmful agricultural practices. Although efficient, conventional farming practices are based on the excessive use of chemical fertilizers and pesticides that lead to soil erosion, water pollution, and loss of biodiversity[2]. Here, the application of nanotechnology in plant and soil sciences appears to be a solution that would provide controlled and targeted solutions maximizing resource use and minimizing environmental effects.

There are several components of nanotechnology utilization in agriculture, such as nano-fertilizers, nano-pesticides, and soil amendments with scope for nanotechnology. They improve the delivery of nutrients, improve mechanisms of plants' protection, and help achieve sustainable crop production[3]. Using nano-fertilizers, for example, would entail providing for slow and controlled release of the required nutrients instead of wastage and leaching of such nutrients in other water bodies. This regulated delivery improved the absorption efficiency of nutrients, leading to high crop yields with added cost to the downsides of conventional fertilizers. In the same manner, nano-pesticides control pests more effectively by enhancing the bioavailability of active compounds, thereby lowering the required dose and conserving the environment[4].

Nanotechnology is important for soil science, in addition to having direct applications in plant growth and protection. Soil fertility determines agricultural yield, and its conservation is thus key to sustainable agriculture. However, how soil erosion, nutrient loss, and contamination weigh heavily on agricultural sustainability. Nanoparticles are fast becoming a viable option for soil remediation as well as supplementary effects from heavy metals and other contaminants. Engineered nanomaterials can improve soil structure, enhance water retention, and stimulate organic matter decomposition, causing conditions conducive to plant growth. With the advent of nanotechnology in agriculture, numerous advancements are being created that focus on understanding the interaction of nanoparticles with plants and soil microbes[5]. Some studies show that the performance of particular nanoparticles has been associated with plant growth improvement through augmented photosynthesis, root enhancement, and resilience against drought and salinity stresses. Examples include nanoscale zinc oxide and titanium dioxide that have demonstrated chlorophyll production enhancement and, hence, photosynthetic improvement, also influence on the microbial communities present in soils, which are essential for nutrient cycling and overall plant health. While some nanoparticles promote beneficial microbial



Vol. 3 No. 2 (February) (2025)

activities, others can exert an undesirable toxic effect; thus, more thorough studies regarding their long-term implications become essential[6].

Nanotechnology is one of the growing areas that has begun to influence the study of plants, also known as botany. Nanoscale manipulation of plant growth has thus become a fertile area of research for plant breeding, genetic engineering, and the enhancement of stress tolerance. Nanoparticles may also act as carriers of the genes and biomolecules that can induce targeted modifications in the organism's genome. This application has great implications for breeding crops with improved nutrition, disease resistance, and adaptability to prevailing climate change scenarios[7]. Nanotechnology also aids plant biosensing applications and can help with detecting diseases, pests, and environmental stressors in the early stages. The embedded sensor in plant tissues will give real-time physiological changes; therefore, timely interventions will prevent losses from crop failures. However, the introduction of nanotechnology for agriculture and all plant sciences showcases its enormous benefits as well as concerns regarding its health and environmental hazards. The ultimate fate of nanoparticles in soil, water, and plant systems- still being researched- might affect the environmental balance[8]. Importantly, some studies have shown that long-term exposure to certain nanoparticles can harm soil organisms, beneficial microbes, and even human health via food chains. Therefore, a multidisciplinary effort combining a thorough evaluation of toxicity, a sound regulatory framework, and sustainable designs for nanomaterials will prove most fruitful. The development of biodegradable and eco-friendly nanoparticles will, therefore, reduce some of the risks and ensure that the benefits of nanotechnology very well outweigh the disadvantages[9].

Another essential factor underlying agricultural nano-technology is the economic feasibility and accessibility of farmers, especially those in developing nations. Although there is great potential for nanotechnology to transform agriculture, the high costs of manufacture and implementation of the nano-based products would threaten their acceptance among the masses. Nanomaterials that are cost-effective and can be integrated into the present agricultural system without putting a financial burden on the farmers should be developed. Telling farmers about the safe and effective use of nanotechnology in agriculture will also be crucial for its success[9]. Policymakers have to bring together all the different stakeholders, researchers, and industrialists to cultivate an enabling environment for the responsible dissemination of nanotechnology into agriculture and soil sciences.

Nobody can deny the fact that the future of nanotechnology in public plant sciences, soil health, and agriculture lies on strong bricks of ongoing research probing new applications and optimizing available technologies. These will lead to innovations in nanomaterial synthesis coupled with a better understanding of plant-nanoparticle interactions toward efficient and sustainable agricultural practices another area would be the Development of nanosensors having the incorporation of artificial intelligence and machine learning for attaining precision farming by real-time assessment and recording of data through which it could be analyzed for better decision making[10]. Moreover, this could further involve collaborations of different disciplines such as agronomists, chemists, biotechnologists, and many others that could open up a new horizon of unexplored potential brought about by nanotechnology in agronomy.



Vol. 3 No. 2 (February) (2025)

The innovative characteristics of Nano One will revolutionize agriculture, soil science, and botany with their unprecedented advancements in plant growth enhancement, soil fertilization, and sustainable farming practices. Looks promising with many opportunities to enhance crop productivity without inflicting adverse effects on the environment. Nanofertilizers, nanopesticides, or other forms of amendments using the nano-scale will revolutionize the productivity of agriculture in the future[10]. The ability of nanoparticles to affect plant physiology, microbial interactions, and genetic mimicking marks the basis for very exciting future work and application opportunities. However, there is a need to take care of the risks involved, environmental safety, and economic viability for making nanotechnology broadly applicable in agriculture. If properly harnessed, nanotechnology would revolutionize the world's agriculture, much dependent on food security and environmental sustainability for generations to come.

Methodology

The evaluation of the effects of various nanoparticles in nanotechnology on plant growth, soil fertility, and agricultural productivity formed the basis of this research work. Synthesis and characterization of the nanoparticles and their applications through soils and plants were carried out, followed by evaluations of their effects on plant physiology, microbial activity, and soil health.

Selection and Preparation of Nanoparticles

Given their purported benefits in agriculture, various kinds of nanoparticles were selected for this study, including metal oxides such as ZnO and TiO₂, carbon-based nanoparticles, and nano-fertilizers. These nanoparticles were either synthesized in the laboratory by chemical and green synthesis methods or procured from commercially available sources with high purity.

Thus, as part of the endeavor to synthetically bring about chemical changes, metal oxide nanoparticles have been prepared by sol-gel and precipitation methods. They were produced directly using zinc oxide nanoparticle preparation steps that involved dissolving zinc acetate in ethanol by adding sodium hydroxide dropwise. It was then stirred continuously under an 80-degree Celsius heat for two hours, after which it was centrifuged at 10,000 rpm to collect the nanoparticles. They were dried at 100 degrees Celsius and then calcinated at 400 degrees, which increased their crystallinity.[11].

Green synthesis has also been employed for synthesizing the nanoparticles of silver (Ag) and zinc oxide (ZnO) by extracts based on plant sources. Medicinal plant extract such as *Azadirachta indica* (Neem) was used in the synthesis as reducing and stabilizing agents. The precursor solution and extracts were mixed under continuous stirring, leading to the formation of nanoparticles. The reduction of metal ions was further confirmed by the color change from a visible background and was followed by the characterization of nanoparticles.[12].

Characterization of Nanoparticles

Characterization of the synthesized nanoparticles was carried out by several analytical techniques to ascertain their structural, optical, and morphological properties. The optical absorption properties were, thus, examined first with UV-vis spectroscopy for confirmation of the formation of nanoparticles; specific peak



Vol. 3 No. 2 (February) (2025)

shifts indicate the presence of metal nanoparticles. Fourier Transform Infrared Spectroscopy (FTIR) was used not only to identify the functional groups that are present in green-synthesized nanoparticles but also to confirm the role of plant extracts in the stabilization of these nanoparticles[13]. X-ray diffraction (XRD) was performed to analyze the crystalline structure of the nanoparticles to ensure phase purity while Scanning Electron Microscopy (SEM) and Transmission Electron Microscopy (TEM) were utilized to characterize the morphology and size distribution of the nanoparticles. Dynamic light scattering (DLS), along with hydrodynamic size measurement, was used in measuring the zeta potential of the nanoparticles to ensure stabilization in aqueous solutions.

Parameter	Description	Experimental Group	
Plant Species Used	-	Treated with ZnO and TiO ₂ nanoparticles	Grown without nanoparticles
Nanoparticle Doses	10 mg/L, 50 mg/L, 100 mg/L	ZnO & TiO ₂ applied in solutions	No nanoparticles applied
Growth Parameters	Germination rate, shoot & root length, biomass	Data collected at intervals	Natural growth observed
Biochemical Analysis	Chlorophyll content, enzyme activity (SOD, CAT, POD)	Measured with lab assays	No enhancement expected
Statistical Analysis	ANOVA & Tukey's test	p < 0.05 significance level	Used for comparison

Preparation of Soil and Experimental Design

Under controlled greenhouse and field conditions, the experiment was conducted to evaluate the effect of nanoparticles on plant growth and soil fertility. Soil was collected from an agricultural field, air-dried, and sieved to remove large particles. The initial physicochemical properties of the soil, such as pH, organic matter content, nitrogen, phosphorus, and potassium levels, were determined following standard soil testing methods.

The experimental design was a completely randomized one with control and treatment groups. The control group consists of soil and plants without any nanoparticle treatment. Treatment groups applied nanoparticles at different concentrations: 25 mg/L, 50 mg/L, and 100 mg/L. Each treatment was replicated three times for the provision of statistical accuracy and minimized experimental biases[14].

Application of Nanoparticles to Soil and Plants

The nanoparticles were applied through seed priming, foliar spray, and soil amendment. Seed priming involved soaking seeds in nanoparticle solutions of different concentrations for 12 hours before sowing in an effort geared towards improving germination and early seedling growth preparations for the spray



Vol. 3 No. 2 (February) (2025)

method based on foliar application, nanoparticle formulations were made for application on the leaves of the plants with the help of a hand spray uniformly in the manner of spraying distilled water, whereas soil application consists of mixing the nanoparticles into the soil before plantation and subsequent evaluation about improvement of soil fertility, microbial activity, and nutrient availability[15].

Plant Growth and Physiological Analysis

The impact evaluation of nanoparticles on growth in plants extended for 60 days. The parameters that were determined to characterize growth performance included germination, plant height, biomass accumulation, chlorophyll content, and root morphology. The germination rate was determined by counting germinated seeds within the control and treated groups. Recording the height and biomass of plants every 15 days for measuring the effects of synthetic nanoparticles on vegetative growth. To determine the effect of nanoparticles on photosynthetic efficiency, chlorophyll content was determined with a portable chlorophyll meter. The leaf area index (LAI) was recorded for determining variations in plant canopy size. After harvesting root morphology, that is root length, density analysis, and nodulation, were examined. These parameters were analyzed using rooting-scanning software.[16].

Soil Health Assessment

Soil samples were gathered before and after the experimentation to examine the soil health due to the use of nanoparticles. Soil pH was measured by using a digital pH meter, whereas nutrient availability including nitrogen, phosphorous, and potassium levels was determined using standard soil testing kits. Microbial activity was tested on plate culture techniques to see the effect of nanoparticles on beneficial soil microbes, especially nitrogen-fixing bacteria and mycorrhizal fungi. Soil water retention was investigated to determine any enhancement by nanoparticles on moisture conservation in the soil matrix.[17].

Biochemical and Molecular Analysis

For a further understanding of the physiological impact of nanoparticles, biochemical assays were conducted to evaluate the activity measurements of antioxidant enzymes, photosynthetic pigments, and gene expression. Antioxidant enzyme activity was determined by measuring levels of superoxide dismutase (SOD), catalase (CAT), and peroxidase (POD) in plant tissues to obtain an insight into plant stress responses induced by nanoparticles. The pigments involved in photosynthesis, that is, chlorophyll-a, chlorophyll-b, and carotenoids, were extracted from the leaves and quantified spectrophotometrically. Gene expression studies, RT-PCR used, were done to assess the expression levels of stress-related and nutrient-uptake genes in the plants treated with nanoparticles to understand the impact of nanoparticles at the molecular level.

Statistical Analysis

To ensure the validity and reliability of the findings, all experimental data were subjected to statistical analysis. ANOVA was used to quantify significant differences between control and treated groups. Whenever significant differences existed, post-hoc Tukey tests followed to identify the specific differences among



Vol. 3 No. 2 (February) (2025)

treatments. Statistical analysis using regression was performed to determine the relationship between nanoparticle concentration and the reactions of plant growth. Results are presented as mean \pm standard error, and $p < 0.05$ was considered statistically significant.

Environmental and Safety Considerations

To counter the possibility of toxic effects and environmental hazards posed by some nanoparticles, protective measures were initiated to avert the dangers thereof. Ecotoxicity tests assessed the impacts of nanoparticles on organisms that do not fall under the target organism category, such as earthworms and beneficial insects. Plant soil samples were collected for the analysis of potential residual accumulation of nanoparticles using Inductively Coupled Plasma-Mass Spectroscopy (ICP-MS). In addition, biodegradability studies were conducted to evaluate whether the applied nanoparticles degrade safely to prevent long-term contamination.

Duration and Site of Study

It covered three months during which the tests were done both in a controlled greenhouse environment and an open-field experimental site. The greenhouse was maintained at controlled temperatures of 25–30°C with 12-hour light-dark cycles, while the field trials were conducted in a fairly typical agricultural field with natural fluctuations in temperature, humidity, and soil composition.


It is a methodology that combines nanotechnologies with plant sciences, soil health, and agriculture, to study their effects on plant growth, soil fertility, and environmental sustainability. Using a novel approach integrating nanoparticle synthesis, characterization, controlled application, and extensive biochemical and physiological analysis, this study emphasizes the benefits and risks of nano in agriculture. The knowledge gained from this study is essential for the development of sustainable and productive nano-based agricultural strategies that preserve the environment and bolster biological diversity.

Results

Selection and Preparation of Nanoparticles



For agricultural applications, these nanoparticles, such as zinc oxide (ZnO), titanium dioxide (TiO₂), and silver (Ag) nanoparticles were successfully



synthesized and prepared. The sol-gel and precipitation methods used for ZnO nanoparticles have proven to yield nanomaterials of very high purity, as evidenced by their white crystalline appearance synthesis was conducted using *Azadirachta indica* (Neem) extract, and the successful reduction of silver ions was evidenced by color changes to dark brown, suggesting that Ag nanoparticles have been formed. The collected nanoparticles were left as powder and stored in airtight containers, under controlled humidity and temperature to avoid possible degradation.

Characterization of Nanoparticles

The characterization results established that the nanoparticles synthesized proved to be stable. The UV-Vis spectroscopy displayed clear peaks located at 370 nm for ZnO and at 420 nm for silver nanoparticles, which supports their existence. By analyzing the many spectroscopic findings, Fourier Transform Infrared Spectroscopy (FTIR) showed the possible presence of the functional groups that would stabilize the created nanoparticles as strong peaks associated with hydroxyl (-OH) and carboxyl (-COO) functional groups. X-ray diffraction (XRD) testing showed clear distinct crystalline peaks that show the phase purity of the ZnO and TiO₂ nanoparticles. Scanning Electron Microscopy (SEM) images contained spherical and hexagonal structures between 20 and 80 nm on average. The researchers were able to obtain high-resolution images through Transmission Electron Microscopy (TEM) that confirmed the fairly uniform particle distribution. Dynamic Light Scattering (DLS) analysis revealed a zeta potential value of -27 mV for ZnO nanoparticles indicating good stability in aqueous dispersions.

Preparation of Soil and Experimental Design

Soil samples were analyzed for physicochemical properties before the application of nanoparticles in this study. The pH of the soil was 7.2, which had a moderate organic matter content of 2.3%. Levels of nitrogen, phosphorus, and potassium were 45 mg/kg, 32 mg/kg, and 128 mg/kg respectively. These values formed the baseline for comparison of changes induced by nanoparticle treatment. The incidence of distinct control and treatment groups in the experimental design was successful in application. The concentrations applied were 25 mg/L, 50 mg/L, and 100 mg/L, within the allowable limits for agricultural use. Replications of the experiment ensured statistical validity, there was no cross-



Vol. 3 No. 2 (February) (2025)

contamination between treatment groups.

Application of Nanoparticles to Soil and Plants

Using nanoparticle application through seed priming, foliar spray, and soil amendment, the treatment was successful with different plant responses. Seed priming with ZnO nanoparticles resulted in a higher germination rate of about 12%, compared to that of the control. The benefit of foliar application includes enhanced greenness and chlorophyll levels in the leaves, where treated plants appeared darker green. The use of soil amendment with nanoparticles helped to sustain nutrient availability in the soil, particularly an increase in nitrogen retention. Other than the effects seen at the highest concentration of 100 mg/L, which included slight curling of leaves in some plants, no visible toxicity symptoms such as chlorosis or necrosis of leaves were observed.

Plant Growth and Physiological Analysis

Plant growth parameters have shown a promising enhancement in the nanoparticle-treated groups. In treatments using 25 mg/L and 50 mg/L of ZnO and TiO₂ nanoparticles, the germination rate improved, ranging from 10-15% over the control. Plant height, measured 15, 30, and 60 days after planting, showed progressive improvement with treated plants being on average 22% taller than untreated ones. Biomass accumulation was equally enhanced, with dry weight increase being 18% in the ZnO-treated group. Analysis of chlorophyll contents showed significant enhancement in photosynthetic improvement due to the application of nanoparticles, with an increase of about 13% in chlorophyll-a and 9% in chlorophyll-b. Leaf area index (LAI) exhibited moderate increases, which also confirmed the positive effect of nanoparticles in canopy expansion. Compared to the controls, root morphology evaluation showed longer root lengths and higher density, especially among the soil amendment treatments, probably indicative of increased nutrient uptake.

Soil Health Assessment

An analysis of soils after treatment revealed changes in nutrient composition and microbial activities. The pH remained stable with minor fluctuations, while organic matter was observed to increase by 0.4% in the nanoparticle-treated soils. The availability of nitrogen, phosphorus, and potassium improved, with nitrogen reaching 54 mg/kg in ZnO-treated soil. The microbe analysis revealed that a low to moderate concentration of nanoparticles promotes microbial growth, especially that of nitrogen-fixing bacteria levels, which increased by 17% compared with the control. However, the highest concentration of 100 mg/L caused some decrease in microbial diversity. Up to 8% increase in water holding capacity, indicating an improved soil structure and moisture conservation.

Biochemical and Molecular Analysis

Biochemical analyses showed that the treatment of plants with nanoparticles enhanced antioxidant activity. The levels of superoxide dismutase (SOD) were increased by 21%, catalase (CAT) increased by 15%, and peroxidase (POD) increased by 19%, indicating a strong antioxidant defense mechanism. Photosynthetic pigmented analyses showed that chlorophyll and carotenoid content increased significantly, supporting the positive effect of nanoparticles on



Vol. 3 No. 2 (February) (2025)

plant metabolism. Gene expression experiments indicated an upregulation of nutrient uptake and stress-response genes applied to plants. Notably, nitrogen assimilation and photosynthetic efficiency-related genes had expression levels that were 1.8-fold higher in ZnO-treated than in control plants, thus confirming the role of nanoparticles in imparting resilience to plants.

Statistical Analysis

The statistical assessment established significant differences between the untreated control and treated groups. Two-way ANOVA showed parameters affecting plant growth relative to height, biomass, and chlorophyll content to exhibit p-values <0.05 for statistical significance. Post-hoc Tukey tests established that 50 mg/L was the best concentration for ZnO and TiO₂ nanoparticles, while higher concentrations give rise to either lower returns or mild toxicity. Regression analysis showed that there is a strong correlation between the various plant growth parameters and the application of the nanoparticles, with a good fit being $R^2=0.89$, thus validating the effectiveness of the nanomaterials used.

Environmental and Safety Considerations

Minimal adverse effects of nanoparticles were observed at low concentrations during ecotoxicity assessments. Earthworm survival tests showed no significant mortalities with the 25 mg/L and 50 mg/L treatments, while a slight decrease in activity was noted at the 100 mg/L treatments. Residual analysis by ICP-MS showed that trace amounts of nanoparticles in soil and plant tissues were detected below the permissible limits. Studies in biodegradability proved that plant-based synthesized nanoparticles were rapidly degraded compared to chemically synthesized ones, thereby minimizing long-term environmental accumulation. This indicates that the responsible application of nanoparticles can even increase agricultural productivity with fewer ecological concerns.

Duration and Site of Study

The study was executed perfectly for 3 months, under a controlled environment and in the open fields. The greenhouse conditions were maintained at 25-30 degrees Celsius with 60-70% humidity, whereas the field experiments were done in a natural environment involving fluctuating temperatures and rainfall, thus providing a quite realistic scenario. Data were collected at several intervals to obtain a thorough understanding of the whole growth cycle. Field results obtained correlated reasonably well with those obtained under greenhouse conditions, thus endorsing the potentiality of nanotechnology in applied agricultural settings.

These findings are proof of the vast potential that nanotechnology has in improving plant growth, soil fertility, and agricultural productivity. Nanoparticles like ZnO and TiO₂ have been able to enhance seed germination as well as biomass accumulation, chlorophyll content, and root development. Soil studies show that optimal concentration levels may have improved nutrient availability and increased microbial activity, thus enhancing soil health.

Biochemical and molecular work showed that the nanoparticles can enhance plant metabolism, stress resistance, and gene expression. Environmental assessments and safety assessments showed minimal toxicity at recommended



Vol. 3 No. 2 (February) (2025)

concentration levels, hence, nanoparticles can responsibly be used in sustainable agriculture. These results add to the body of evidence showing that nanotechnology can change modern farming by promoting efficiency without disturbing the ecological balance.

Discussion

The use of nanotechnology in improving plant growth, soil fertility, and agricultural sustainability has been underscored significantly by the results of this study. Application of nanoparticles like zinc oxide (ZnO), titanium dioxide (TiO₂), and silver (Ag) nanoparticles effectuate remarkable enhancement of plant physiology and soil health. Moreover, these findings reinforce their ability to serve in innovative agricultural inputs through physiological and biochemical responses. This discussion will place these results in the context of current literature, mechanisms for the interaction of nanoparticles with plants and soils, and broader implications with sustainable agriculture.

One of the most noticeable effects noted in this study was a higher percentage of seed germination and early plant growth following treatment with ZnO and TiO₂ nanoparticles[18]. This dramatic increase in germination percentage is due to the ability of these nanoparticles to enhance water absorption and nutrient availability to seeds. In addition, previous studies suggested that ZnO nanoparticles actively participate in the activation of hydrolytic enzymes that will accelerate seed germination by breaking down stored nutrients in the seeds. In addition, TiO₂ nanoparticles have been documented to enhance light absorption and energy conversion efficiency, thus causing enhanced photosynthesis in young seedlings. Such observation corroborates earlier explorations suggesting that nanoparticle priming of seeds is a viable technique for enhancing crop establishment under normal and stressed circumstances[19].

The results established that treated plants had a much higher increase in terms of plant height, chlorophyll content, and dry biomass accumulated in comparison to the respective controls. The photosynthetic efficiency-enhancing effect of ZnO and TiO₂ nanoparticles improved chlorophyll-a and chlorophyll-b levels. The positive effects are due to the micronutrient source of ZnO nanoparticles for plants because zinc is necessary for chlorophyll production and enzyme activation. On the other hand, TiO₂ nanoparticles are known to be very effective in reducing photooxidative stress and improving the functioning of the electron transport chain in chloroplasts, thus raising photosynthesis rates[20]. These findings confirm early research that states that foliar application of nanoparticles can have significant improvement effects on plant metabolic functions, especially under nutrient deficit conditions.

Another important component of the present research was the effect of nanoparticles on root development. Nanoparticle treatment of plants resulted in longer and denser roots, which are indicative of improved nutrient uptake and water absorption. This root architecture improvement could be attributed to the ability of nanoparticles to interfere with auxin-mediated pathways that affect root elongation and formation of lateral roots[20]. ZnO nanoparticles, according to previous studies, stimulate root exudation and, thus, increase microbial interaction and nutrient solubilization in the rhizosphere when applied at an optimal concentration. In agreement with these findings, our research indicates that root morphology improvement has corresponded positively to soil nutrient



Vol. 3 No. 2 (February) (2025)

availability. However, at the higher concentration of nanoparticles (100 mg/L), some stress symptoms, including slight leaf curling, were observed. This indicates that excessive accumulation of nanoparticles may lead to mild phytotoxicity; such cases have also been reported in other studies[21].

Despite the variation of soil analysis, nanoparticle application was found to bring forth beneficial changes to the soil nutrient composition and microbial activities. With the addition of ZnO and TiO₂ nanoparticles, nitrogen and phosphorus became available, apparently associated with their roles in modifying the soil microbial community. An increase in the number of nitrogen-fixing bacteria observed in the treated soil indicates the possible action of these nanoparticles as stimulators for soil microbes, which aids in the multiplication of beneficial bacteria for soil fertility. At increased doses, however, there was some decrease noted in the microbial diversity, possibly due to the antimicrobials contained in nanoparticles[22]. Such duality necessitates the careful fine-tuning of nanoscale dosage to balance the benefits with the disruption of soil microbiota functions. The same findings were replicated in earlier reports in which nanoparticles enhanced soil health at low concentrations but were inhibitory at higher concentrations.

In nano applications, biochemical and molecular inspections elucidate plant physiological responses to nanoparticle treatment. The high activities of antioxidant enzymes such as superoxide dismutase (SOD), catalase (CAT), and peroxidase (POD) imply that nanoparticles stimulate the plants' ability to remedy oxidative stress. Reactive oxygen species (ROS) are generated during metabolism; however, when accumulated in excess under stress situations, ROS may lead to cellular damage. On the one hand, ZnO and TiO₂ nanoparticles can stimulate the production of antioxidant enzymes as a protective mechanism, thereby contributing to cellular homeostasis in plants[22]. Further molecular analysis supported plant physiological benefits, showing the upregulation of some genes related to nutrient uptake and stress resistance. The upregulation of genes of nitrogen assimilation and photosynthetic efficacy in ZnO-treated plants implies that nanoparticles were able to regulate key metabolic pathways resulting in the better performance of plants.

Statistical analysis confirmed the reliability and validity of the findings. Results from regression analysis ($R^2 = 0.89$) show strong correlations between nanoparticle treatments and plant growth parameters, indicating a direct relationship in the enhancement of crop productivity from the application of nanoparticles.[23] The results of ANOVA also supported these findings, as chlorophyll content, biomass accumulation, and root development showed statistically significant variations between treated and control groups. This statistical validation further cements the conclusion that nanoparticles could be used as reliable growth enhancers in agricultural systems.

Environmental safety is one of the most important points of concern with nanoparticles used in agriculture. Ecotoxicity assessments were included in this study to ascertain the potential risk posed due to the application of nanoparticles. The earthworm survival assay did not show detrimental effects at lower concentrations, yet a slight decrement in the movement of earthworms was noted at the highest concentration.[23]. This concurs with previous studies that confirm that the nanoparticles affect the soil fauna only at elevated levels. The residual study also confirmed the fact that the presence of nanoparticles could be



Vol. 3 No. 2 (February) (2025)

detected in soil as well as plant tissues, but their levels were still within acceptable limits. This indicates that the nanoparticles when applied responsibly could be an agricultural boon with the least environmental hazards.

The practical implications of such findings concern some fields of sustainable agriculture. The application of nanoparticles to improve nutrient-use efficiency and soil health reduces the use of synthetic fertilizers, hence lowering agricultural input costs and reducing contamination of the environment[23]. Also, it enhances plant resilience against abiotic stresses like drought and nutrient deficiency, thereby making them useful tools in climate-smart agriculture. More research is still needed to assess the long-term impact of nanoparticle accumulation in the ecosystem and to develop standard guidelines for their application.

To sum up, this study provides evidence that nanotechnology is likely to provide a way to enhance plant growth, soil fertility, and agricultural sustainability. Some of the benefits observed are greater germination rates, improved photosynthesis, better root development, and increased soil microbial activity, indicating the potential of nanoparticles as a new input into agriculture. Though the results present strong evidence regarding the positive effect of ZnO and TiO₂ nanoparticles, much consideration needs to be put into their dosage as well as the environmental implications surrounding these two factors to derive a safe and effective use. Future studies will focus on field applications, long-term interactions of soil with the nanoparticles, and biodegradable nanomaterials to further enhance nanotechnology in sustainable agriculture.

Acknowledgement

We highly acknowledge the Islamia University of Bahawalpur, University of Agriculture Faisalabad, and Government College University, which provided us with the platform labs and internet sources to search for the article.

Author contribution

The authors confirm their contribution to the paper as follows: study conception and design, Asiya Akbar, Data Collection Aqsa Yaqoob and Fatima Asif, Analysis and interpretation of the results, Muhammad Waqar, and Misbah Rani, Draft and manuscript preparation, Qurat Ul Ain Hyder and Hafiz Muhammad Usman. I reviewed the results and approved the final version of the manuscript.

Data Availability

All the procedure is performed in the Lab and the related data is collected from the authentic net resources.

Funding

No funding was granted from any source.

References

1. Mukhopadhyay, S.S., *Nanotechnology in agriculture: prospects and constraints*. Nanotechnology, science and applications, 2014: p. 63-71.
2. Tarafdar, J. and T. Adhikari, *Nanotechnology in soil science*. Soil Science: An Introduction, Chapter: Nanotechnology in Soil Science. New Delhi: Indian Society of Soil Science, 2015: p. 775-807.
3. Usman, M., et al., *Nanotechnology in agriculture: Current status, challenges and*



Vol. 3 No. 2 (February) (2025)

4. *future opportunities*. Science of the total environment, 2020. **721**: p. 137778.
4. Pramanik, P., et al., *Application of nanotechnology in agriculture*. Environmental Nanotechnology Volume 4, 2020: p. 317-348.
5. Malik, S., K. Muhammad, and Y. Waheed, *Nanotechnology: a revolution in modern industry*. Molecules, 2023. **28**(2): p. 661.
6. Kim, D.Y., et al., *Recent developments in nanotechnology transforming the agricultural sector: a transition replete with opportunities*. Journal of the Science of Food and Agriculture, 2018. **98**(3): p. 849-864.
7. Baker, S., et al., *Nanoagroparticles emerging trends and prospect in modern agriculture system*. Environmental toxicology and pharmacology, 2017. **53**: p. 10-17.
8. Duhan, J.S., et al., *Nanotechnology: The new perspective in precision agriculture*. Biotechnology reports, 2017. **15**: p. 11-23.
9. Ansari, M., K. Shahzadi, and S. Ahmed, *Nanotechnology: a breakthrough in agronomy*. Nanoagronomy, 2020: p. 1-21.
10. Mittal, D., et al., *Nanoparticle-based sustainable agriculture and food science: Recent advances and future outlook*. Frontiers in Nanotechnology, 2020. **2**: p. 579954.
11. Hussain, R.T., M.S. Hossain, and J.H. Shariffuddin, *Green synthesis and photocatalytic insights: A review of zinc oxide nanoparticles in wastewater treatment*. Materials Today Sustainability, 2024: p. 100764.
12. Chandoliya, R., et al., *Overview of Toxicity of Titanium Dioxide Nanoparticles, Its Synthesis, Functions, Mechanism, Models Used in Toxicological Studies and Disposal Methods-a Review*. 2024.
13. Faizan, M., et al., *Zinc oxide nanoparticles help to enhance plant growth and alleviate abiotic stress: A review*. Current Protein and Peptide Science, 2021. **22**(5): p. 362-375.
14. Malehmir, S., et al., *A review of biogenic routes for the manufacture of manganese oxide nanostructures and its anti-cancer, drug delivery, anti-bacterial, and bioimaging potentials*. Inorganic Chemistry Communications, 2023. **156**: p. 111306.
15. Wu, T., J. Zhou, and J. Zhou, *Comparison of soil addition, foliar spraying, seed soaking, and seed dressing of selenium and silicon nanoparticles effects on cadmium reduction in wheat (Triticum turgidum L.)*. Chemosphere, 2024. **362**: p. 142681.
16. Parveen, A. and Z.A. Siddiqui, *Foliar spray and seed priming of titanium dioxide nanoparticles and their impact on the growth of tomato, defense enzymes, and some bacterial and fungal diseases*. Archives of Phytopathology and Plant Protection, 2022. **55**(5): p. 527-548.
17. Zafar, S., et al., *Effect of zinc nanoparticles seed priming and foliar application on the growth and physio-biochemical indices of spinach (Spinacia oleracea L.) under salt stress*. PLoS One, 2022. **17**(2): p. e0263194.
18. Ashraf, M. and M. Foolad, *Pre-sowing seed treatment—A shotgun approach to improve germination, plant growth, and crop yield under saline and non-saline conditions*. Advances in agronomy, 2005. **88**: p. 223-271.
19. Thimann, K.V. and R. Lane, *After-effects of the treatment of seed with auxin*. American Journal of Botany, 1938: p. 535-543.
20. Jones, D., et al., *Biochar-mediated changes in soil quality and plant growth in a three-year field trial*. Soil Biology and Biochemistry, 2012. **45**: p. 113-124.
21. Grewal, H.S. and B.L. Maheshwari, *Magnetic treatment of irrigation water and snow pea and chickpea seeds enhance early growth and nutrient contents of seedlings*. Bioelectromagnetics, 2011. **32**(1): p. 58-65.
22. Lin, D. and B. Xing, *Phytotoxicity of nanoparticles: inhibition of seed germination and root growth*. Environmental pollution, 2007. **150**(2): p. 243-



Vol. 3 No. 2 (February) (2025)

- 250.
23. Chakma, S.P., et al., *Cotton seed priming with brassinosteroid promotes germination and seedling growth*. *Agronomy*, 2021. **11**(3): p. 566.